

AD-A163 485


ISOLATION AND CHARACTERIZATION OF AN  
ORGANO Gallium-ARSENIC CLUSTER CRYST (U) DUKE UNIV  
DURHAM NC DEPT OF CHEMISTRY R L WELLS ET AL 18 JAN 86  
DU/DC/TR-01 N00014-83-K-0572 F/G 20/2

1/1

UNCLASSIFIED

NL





MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

12

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER DU/DC/TR-01	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Isolation and Characterization of an Organo- gallium-Arsenic Cluster: Crystal Structure of [(PhAsH)(R <sub>2</sub> Ga)(PhAs) <sub>6</sub> (RGa) <sub>4</sub> ] (R = Me <sub>3</sub> SiCH <sub>2</sub> )		5. TYPE OF REPORT & PERIOD COVERED Technical Report 1985
7. AUTHOR(s) R. L. Wells, A. P. Purdy, A. T. McPhail, and C. G. Pitt		6. PERFORMING ORG. REPORT NUMBER TR-01
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry Duke University Durham, NC 27706		8. CONTRACT OR GRANT NUMBER(s) N00014-83-K0572
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 053-341
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE Jan. 10, 1986
		13. NUMBER OF PAGES 8
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release: distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)  Prepared for publication in <u>J. Chem. Soc., Chem. Commun.</u>		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Organogallium-arsenic Ga-As cluster		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) [(PhAsH)(R <sub>2</sub> Ga)(PhAs) <sub>6</sub> (RGa) <sub>4</sub> ] (R = Me <sub>3</sub> SiCH <sub>2</sub> ) has been isolated as a product of the reaction of PhAsH <sub>2</sub> with (Me <sub>3</sub> SiCH <sub>2</sub> ) <sub>3</sub> Ga and determined by x-ray crystallographic analysis to be a cluster containing an As <sub>7</sub> Ga <sub>5</sub> core.		

AD-A163 485

DTIC FILE COPY

DTIC  
ELECTE  
JAN 31 1986  
S D

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

S N 0102-LF-014-6601

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

86 1 30 007

OFFICE OF NAVAL RESEARCH

Contract N00014-83-K0572

Task No. NR 053-841

TECHNICAL REPORT \* J/DC/TR-01

Isolation and Characterization of an Organogallium-Arsenic  
Cluster: Crystal Structure of  $[(\text{PhAsH})(\text{R}_2\text{Ga})(\text{PhAs})_6(\text{RGa})_4]$   
( $\text{R} = \text{Me}_3\text{SiCH}_2$ )

by

R. L. Wells, A. P. Purdy, A. T. McPhail, and C. G. Pitt

Prepared for Publication

in the

Journal of the Chemical Society, Chemical Communications

Duke University  
Department of Chemistry  
Durham, NC 27706

January, 1986

Reproduction in whole or in part is  
permitted for any purpose of the United States Government

This document has been approved for public  
release and sale; its distribution is unlimited

Accession For	
NTIS	CRA&I <input checked="" type="checkbox"/>
DTIC	TAB <input type="checkbox"/>
Unannounced <input type="checkbox"/>	
Justification	
By	
Distribution	
Availability Codes	
Dist	Avail and/or Special
A-1	<input checked="" type="checkbox"/>



Isolation and Characterization of an Organogallium-arsenic  
Cluster: Crystal Structure of  $[(\text{PhAsH})(\text{R}_2\text{Ga})(\text{PhAs})_6(\text{RGa})_4]$   
( $\text{R} = \text{Me}_3\text{SiCH}_2$ )

Richard L. Wells\*, Andrew P. Purdy, Andrew T. McPhail, and  
Colin G. Pitt\*

Department of Chemistry, Paul M. Gross Chemical Laboratory, Duke University,  
Durham, NC 27706 U.S.A.

$[(\text{PhAsH})(\text{R}_2\text{Ga})(\text{PhAs})_6(\text{RGa})_4]$  ( $\text{R} = \text{Me}_3\text{SiCH}_2$ ) has been isolated as a product  
of the reaction of  $\text{PhAsH}_2$  with  $(\text{Me}_3\text{SiCH}_2)_3\text{Ga}$  and determined by x-ray  
crystallographic analysis to be a cluster containing an  $\text{As}_7\text{Ga}_5$  core.

As part of our studies of synthetic methods and the identification of  
substituents which will permit the isolation of monomeric and oligomeric  
gallium-arsenic compounds, we investigated the reaction of  $\text{PhAsH}_2$  with  
 $(\text{Me}_3\text{SiCH}_2)_3\text{Ga}$ . Primarily, it was of interest to ascertain if steric bulk  
would permit the isolation of compounds of the type  $(\text{PhAsGaCH}_2\text{SiMe}_3)_n$  ( $n =$   
1-4) and suppress formation of polymers similar to the nonvolatile polymeric  
material reported to be the primary product of the reaction of  $\text{PhAsH}_2$  with  
 $\text{Me}_3\text{Ga}$ . Although no evidence for any of the desired compounds was obtained  
when  $\text{PhAsH}_2$  was allowed to react with  $(\text{Me}_3\text{SiCH}_2)_3\text{Ga}$ , the novel cluster  
 $[(\text{PhAsH})(\text{R}_2\text{Ga})(\text{PhAs})_6(\text{RGa})_4]$  ( $\text{R} = \text{Me}_3\text{SiCH}_2$ ), (1), containing an  $\text{As}_7\text{Ga}_5$  core

was eventually isolated from the complex mixture of products formed. In addition, the known compound  $(\text{PhAs})_6^3$  was isolated.

A mixture of  $\text{PhAsH}_2$  and  $(\text{Me}_3\text{SiCH}_2)_3\text{Ga}$  (1.0 : 0.95 mol ratio), warmed in stages to  $50^\circ\text{C}$ , reacted smoothly, as evidenced by the formation of  $\text{Me}_4\text{Si}$  and a non-condensable gas (presumably  $\text{H}_2$ ). The pentane extract of the nonvolatile products afforded a viscous liquid-solid mixture. Crystals of (1)<sup>†</sup> precipitated slowly from the ligroin soluble fraction of this mixture. Crystallization of the pentane-insoluble products from benzene and then tetrahydrofuran afforded  $(\text{PhAs})_6^+$ .

Crystal Data.  $\text{C}_{66}\text{H}_{102}\text{As}_7\text{Ga}_5\text{Si}_6$ , (1),  $M = 1937.12$ , Triclinic, space group  $\overline{P}1$ ,  $a = 15.398(2)$ ,  $b = 21.755(1)$ ,  $c = 15.155(1)$  Å,  $\alpha = 96.43(1)$ ,  $\beta = 114.42(1)$ ,  $\gamma = 100.43(1)^\circ$ ,  $V = 4446.1$  Å<sup>3</sup>,  $Z = 2$ ,  $D_c = 1.447$  g cm<sup>-3</sup>,  $\mu(\text{Cu-K}\alpha) = 56.6$  cm<sup>-1</sup>. The crystal structure was solved by direct methods.<sup>4</sup> Least-squares refinement of atomic positional and thermal (anisotropic As, Ga, Si; isotropic C) parameters<sup>5</sup> converged to  $R = 0.074$  over 12276 absorption-corrected reflections [ $I > 3.0\sigma(I)$ ] recorded on an Enraf-Nonius CAD-4 diffractometer (Cu-K $\alpha$  radiation, incident-beam graphite monochromator;  $\omega$ - $2\theta$  scans,  $\theta_{\text{max}} = 67^\circ$ ). The final difference Fourier map shows residual electron density around the center of symmetry due to partial occupancy by hydrocarbons from the ligroin used to crystallize (1). The calculated density, given above, omits the unknown extent of solvent present.

The structure of one enantiomer and its Ga-As core are presented in Figure 1. The Ga-As core of the cluster may be viewed in several different ways, one of which is to consider it as being comprised of an eight-membered

As(2)-Ga(5)-As(3)-Ga(7)-As(9)-Ga(10)-As(11)-Ga(12) ring in a distorted crown form with three axial Ga sites linked on one side of the ring by an As(1)-As(8) unit, while on the opposite side of the ring three axial As sites are bridged by an As(4)-Ga(6) moiety. Alternatively, it may be viewed as being comprised of two nonplanar five-membered rings [(As(4)-As(2)-Ga(12)-As(11)-Ga(6) and As(8)-As(1)-Ga(7)-As(3)-Ga(5)] linked by As(1)-Ga(12), As(2)-Ga(5), and As(3)-Ga(6) bonds, and bridged by an As(9)-Ga(10) unit. The single hydrogen atom bonded directly to an arsenic atom could not be located unequivocally in a difference Fourier synthesis, but of the three possible sites, viz. As(4), As(8), or As(9), the latter seems most likely since this atom is adjacent to the unique gallium atom bearing two  $\text{Me}_3\text{SiCH}_2$  groups, i.e. Ga(10). Support for this assignment derives from the fact that the mean bond angles at As(4) and As(8) are  $97.1$  and  $97.8^\circ$ , respectively; whereas at As(9) the mean is  $113.3^\circ$ . Ga-As bond lengths in (1) range from  $2.450(1)$  to  $2.553(1)$  Å, with the mean at  $2.503$  Å being intermediate between the means of  $2.524$  Å in the centrosymmetric dimer  $[(\text{Me}_3\text{SiCH}_2)_2\text{AsGaPh}_2]_2$ <sup>5</sup> and  $2.492$  Å in monomeric  $(\text{Mes}_2\text{As})_3\text{Ga}$ <sup>6</sup> which contains a trigonal planar gallium atom.

To the best of our knowledge, compound (1) is the first cluster compound containing a core of gallium and arsenic atoms to be isolated and characterized. Although it is not possible to account precisely for its formation, we have previously observed that hydrogen and diarsines are by-products of the reactions of hindered gallanes, e.g.  $(\text{Me}_3\text{SiCH}_2)_3\text{Ga}$ , and secondary arsines.<sup>7</sup> Thus, it is not too surprising to find that a compound

such as  $(\text{PhAs})_6$  is produced in the reaction of  $\text{PhAsH}_2$  with  $(\text{Me}_3\text{SiCH}_2)_3\text{Ga}$ , and that (1) contains As-As bonds as well as Ga-As bonds

We thank the Office of Naval Research for financial support.



## References

- 1 O. T. Beachley and R. G. Simmons, Inorg. Chem., 1980, 19, 1021.
- 2 O. T. Beachley and G. E. Coates, J. Chem. Soc., 1965, 3241.
- 3 K. Hedberg, E. W. Hughes, and J. Waser, Acta Cryst., 1961, 14, 369.
- 4 MULTAN11/82. Principal author: P. Main, University of York, England.
- 5 R. L. Wells, A. P. Purdy, A. T. McPhail, and C. G. Pitt, Abstracts of Papers, 189th National Meeting of the American Chemical Society, Miami Beach, FL, April, 1985 American Chemical Society; Washington, DC, 1985, INOR 26.
- 6 C. G. Pitt, K. T. Higa, A. T. McPhail, and R.L. Wells, J. Am. Chem. Soc., submitted for publication.
- 7 R. L. Wells, A. P. Purdy, and C. G. Pitt, unpublished results.

## Footnotes

+ Compound (1) mp 153 °C, dec. Anal. Calcd for  $C_{66}H_{102}As_7Ga_5Si_6$ : C, 40.92; H, 5.31. Found: C, 41.27; H, 5.59.  $^1H$  nmr (ppm) (250MHz),  $C_7D_8$ : complex -0.6 to 0.6 ( $Me_3SiCH_2$ ), and 6.7 to 8.4 (Ph); five peaks 3.2 to 4.1 (AsH).

† Identified by comparison of recorded x-ray diffraction data with those given in ref. 3.

§ The atomic co-ordinates for this work are available on request from the Director of the Cambridge Crystallographic Data Centre, University Chemical Laboratory, Lensfield Rd., Cambridge CB2 1EW. Any request should be accompanied by the full literature citation for this communication.

Figure 1. Structure of a. compound (1), and  
b. the Ga-As core of compound (1)





Fig. 1a.



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 413 800 N. Quincy Street Arlington, Virginia 22217	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 5042 Crane, Indiana 47522	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Washington, D.C. 20360	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
DTNSRDC Attn: Dr. G. Bosmajian Applied Chemistry Division Annapolis, Maryland 21401	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1		

END

FILMED

3 - 86

DTIC